arXiv:physics/0312062v1 [physics.bio-ph] 10 Dec 2003

The Inverse Variational Problem and Logistic Self-Regulated Systems

Á. G. Muñoz S.[1,2*], D. Sierra Porta[1,2], T. Soldovieri[1]
[1]*Grupo de Investigaciones de Física Teórica (GIFT) Departamento de Física.*
Facultad de Ciencias. La Universidad del Zulia. Maracaibo 4004, Venezuela. and
[2]*Postgrado en Física Fundamental, Centro de Astrofísica Teórica (CAT),*
Facultad de Ciencias, Universidad de Los Andes. Mérida, 5101, Venezuela.

There exists in nature many examples of systems presenting self-limiting behaviour: population dynamics, structure engineering, Townsend's electron breakdown, nuclear decay in radioactive equilibrium, histeresis process, meteorological models, etcetera. In this work we call attention to the advantages the use of a variational formulation should provide to the study of self-regulated systems, such as a unified description of the related phenomena, further comprehension of the internal structure and symmetries of the related equations, and the determination of the equilibria points via the energy function. We study the case of logistic systems, obtaining explicitly several s-equivalent Lagrangeans corresponding to the Verhulst's equation. Some dynamical properties are discussed in the light of this approach. Expressions for the mean energy function are also obtained.

Keywords: logistic equation, Verhulst's Lagrangean, self-regulated systems.
PACS: 02.30.Zz, 45.20.Jj, 87.23.Cc

I. INTRODUCTION

The Lagrangean formalism is a very powerful tool in physics, presenting in a single formula all the dynamical information of a system. Nevertheless, it is widely accepted that a variational principle cannot be constructed for an arbitrary differential equation [1]: there exists a strict mathematical theorem that shows its existence for a given situation, and whose application reduces the number of equations in physics that have a Lagrangean-Hamiltonian formulation [2, 3]. Due to its importance, efforts have been directed to make several methods to circumvent this condition and to construct modified variational principles [1, 4, 5, 6].

However, there is an interesting different approach, known as the inverse problem of the variational calculus, or IPVC (see, for example, [7, 8]), that has its historical origin in the times of Helmholtz [9]. It consists in studying the existence and uniqueness (or multiplicity) of Lagrangeans for systems of differential equations, meaning finding the Lagrangean, if it exists, from the equations of motion, instead of the traditional approach. This proceedure has become quite useful and one important result is that of Hojman et al. [10], who have proved that it is possible to construct the Lagrangean for any regular mechanical system as linear combination of their own equations of motion. This particular construction, for example, is much wider than the traditional definition $L = T - V$, which is only true when the "forces" involved are derivable from position-dependent potentials (or very few cases of velocity-dependent potentials), therefore it may be used for general non-conservative systems.

Whereas the equations of the models employed to describe auto-regulated phenomena can be understood as equations of motion in the variational sense, the latter approach shows itself to be one of the most adequate to obtain the Lagrangean formulation of the problem. Its application to the study of self-limiting processes may provide additional understanding into the internal structure of these phenomena, also enables the use of a well known mathematical machinery to find conserved quantities, equilibria and stability cases, and other dynamical properties.

In particular, the method proposed by Hojman et al. provides the desired Lagrangean both elegantly and without many complications, relating directly the constants of motion with the problem. Bearing this in mind, in the following pages we treat the problem of self-regulated systems described by the Verhulst's Logistic Equation (VLE, from now on) [11, 12] by means of the Hojman proceedure, showing how a variational formulation can be easily obtained once we know the equation of motion and some additional information about the system. We have chosen the VLE because it had been well studied, thus we can easily compare the results here obtained to evaluate the approach, and also becasuse the model is simple enough as to give an idea, without involving too much time, of the way a self-regulatory system works.

Historically this equation, firstly proposed by P.F.

---

[*] To whom the correspondence should be adressed. E-mail: agmunoz@mail.luz.ve

Verhulst [11] in 1838 as a continuous non-linear population growth model with a self-limiting density dependent mechanism, was suggested as a direct generalization of the Euler and Malthus [13] studies about population dynamics.

If $n(t)$ is the population at a time t, A^{-1} the representative time scale of response of the model to any change in the population and B define the carrying capacity of the enviroment (see below), then the logistic equation can be written as

$$\frac{dn}{dt} = An\left(1 - \frac{n}{B}\right) \tag{1}$$

The solution of equation (1), for $n(0) = n_0$ is:

$$n(t) = \frac{n_0 B e^{At}}{B + n_0\left(e^{At} - 1\right)} \tag{2}$$

and we can inmediatly show that $\lim_{t\to\infty} n(t) = B$ (for all n_0), meaning, of course, that the carrying capacity corresponds to the the maximum size of the stable steady state population. In the Figure 1 the qualitative behavior of (2) is presented. For $n_0 < B$ the profile is the characteristic sigmoid of the model, and for $n_0 > B$ the behavior is similar to an exponentially decay function.

Indeed, Verhulst knew [14] that there were "biological obstacles" in the natural population growth, but to write a general mathematical expression for them was hardly possible, because each obstacle could affect in a different way the growth itself. Instead, he proposed a general "retardatrice" density dependent ($\propto n^2$) function, a kind of compensating effect of overcrowding. However, it is important to remark that, even a very simple model, it is really useful as it matches a wide variety of different contexts, not only in population dynamics [12], but also in biological growth (see for example, [14]), radioactive decay series [15], and even in some reaction-diffusion models [16]. Effectively, all these cases (and more) can be adequately described by means of a characteristic saturation curve like (2) and in these cases an interpretation of the phenomenona by means of change in the population of some element (individuals, cells, isotopes, ...) can be made.

The self-regulated systems, not only those described by the VLE, are present everywhere: certain aspects in stability of structures (firstly pointed out by Euler [17]), Townsend electron breakdown [18], histeresis and magnetization processes [19], the Amdhal law for scalability of computer programs and a long etcetera. As far as these phenomena are studied sometimes by quite different disciplines of science, it is interesting to explore if it is possible to provide a unified description for them in terms of families of Lagrangeans (or Hamiltonians), that also would help to classify the systems by their dynamical properties.

The distribution of this work is the following: Next section contains an abstract of the Hojman et al. method, which we use in Section 3 to construct the Verhulst's Lagrangian and Hamiltonian. Section 4 is devoted to the study of the energies related to the system, its equilibrium points and its constant of motion. Finally the concluding remarks are presented in Section 5.

II. THE HOJMAN ET AL. METHOD

For the purposes of this paper it is useful to work with the usually denominated second order approach [10]. In this context, the equation of motion for a mechanical system arises from a set of m differential equations:

$$G^i \equiv \ddot{q}^i - F^i\left(q^j, \dot{q}^j, t\right) = 0, \qquad i, j = 1, ..., m \tag{3}$$

where the q^i are the generalized coordinates and the point means total temporal derivative.

In the IPVC the Lagrangean $L\left(\dot{q}^j, q^j, t\right)$ is constructed such that relations (3) can be effectively deduced via the Euler-Lagrange equations. The existence of such a Lagrangean is studied by means of the nowdays called Helmholtz conditions [20]

$$\frac{\partial G_i}{\partial \ddot{q}^j} = \frac{\partial G_j}{\partial \ddot{q}^i} \tag{4}$$

$$\frac{\partial G_i}{\partial \dot{q}^j} + \frac{\partial G_j}{\partial \dot{q}^i} = \frac{d}{dt}\left(\frac{\partial G_i}{\partial \ddot{q}^j} + \frac{\partial G_j}{\partial \ddot{q}^i}\right) \tag{5}$$

$$\frac{\partial G_i}{\partial q^j} - \frac{\partial G_j}{\partial q^i} = \frac{1}{2}\frac{d}{dt}\left(\frac{\partial G_i}{\partial \dot{q}^j} - \frac{\partial G_j}{\partial \dot{q}^i}\right) \tag{6}$$

Nevertheless, these conditions do not give any warranty about uniqueness. Two Lagrangeans are said to be solution-equivalents (or s-equivalents) if they just differ by a global multiplicative constant, η, and a total time derivative of some gauge $\Lambda\left(\dot{q}^j, q^j, t\right)$:

$$\eta L = \widetilde{L} + \frac{d\Lambda}{dt} \tag{7}$$

The different systems of equations they provide, however, have exactly the same equations of motion.

We can proceed now to describe briefly the method. It enables us to write $\widetilde{L}$ as a linear combination of the known equations of motion; then for $i, j = 1...m$ (the degrees of freedom),

$$\widetilde{L} = \mu_i \left[\ddot{q}^i - F^i \left(q^j, \dot{q}^j, t \right) \right] \tag{8}$$

where

$$\begin{aligned} \mu_i \left(q^j, \dot{q}^j, t \right) &\equiv D_1 \frac{\partial D_2}{\partial \dot{q}^i} + ... + D_{2m-1} \frac{\partial D_{2m}}{\partial \dot{q}^i} \qquad (9) \\ &= -\frac{\partial \Lambda}{\partial \dot{q}^i} \qquad (10) \end{aligned}$$

and the D_{2m} are constants of motion of the mechanical system and the D_{2m-1} are arbitrary functions whose arguments are constants of motion. One possible form for the D_{2m-1} functions, given the D_{2m} conserved quantities, is presented in reference [7].

When the conserved quantities are unknown then the problem is reduced to find μ_i such that the following system is satisfied:

$$\begin{cases} \frac{\partial \mu_i}{\partial \dot{q}^j} = \frac{\partial \mu_j}{\partial \dot{q}^i} \\ \frac{\bar{d}}{dt} \left(\frac{\bar{d}}{dt} \mu_i + \mu_j \frac{\partial F^j}{\partial \dot{q}^i} \right) - \mu_j \frac{\partial F^j}{\partial q^i} = 0 \end{cases} \tag{11}$$

and

$$\det \left[\frac{\partial}{\partial \dot{q}^j} \left(\frac{\bar{d}}{dt} \mu_i + \mu_k \frac{\partial F^k}{\partial \dot{q}^j} \right) + \frac{\partial \mu_i}{\partial q^j} \right] \neq 0 \tag{12}$$

where the on-shell derivative $\frac{\bar{d}}{dt}$ is defined as

$$\frac{\bar{d}}{dt} \equiv F^i \frac{\partial}{\partial \dot{q}^i} + \dot{q}^i \frac{\partial}{\partial q^i} + \frac{\partial}{\partial t} \tag{13}$$

For further details the reader is exhorted to review [7, 10] and the references therein. In the following section we make use of this proceedure to obtain the Lagrangean corresponding to the VLE.

III. CONSTRUCTING THE VERHULST'S LAGRANGEAN

For convenience, we shall write (1) as 14

$$\dot{q} \equiv \frac{dq}{dt} = kq \left(B - q \right) \tag{14}$$

with $k \equiv A/B$, and we shall consider the one-dimensional problem.

In our case, equation (3) is

$$\ddot{q} - k\dot{q}(B - 2q) = 0 \tag{15}$$

where $\dot{q}$ is given by (14). Explicitly in terms of the generalized coordinate (on shell), we have

$$\ddot{q} - k^2 q(B - q)(B - 2q) = 0 \tag{16}$$

together with the initial conditions

$$\dot{q}(0) = kq_0 \left(B - q_0 \right) \tag{17}$$

$$q(0) = q_0 \tag{18}$$

Note at this point that, indeed, the Verhulst Lagrangean exist, for equation (16) satisfies trivially the Helmoltz conditions.

The second order Hojman et al. method provides then

$$\widetilde{L_V} = \mu \left[\ddot{q} - k^2 q(B - q)(B - 2q) \right] \tag{19}$$

We just need to determine the factor μ, which in this case can be written as

$$\mu = C_1 \frac{\partial C_2}{\partial \dot{q}} \tag{20}$$

There are several ways to proceed now. For example, observe that from the initial conditions and (16) we can obtain the only constant of motion, C_2, the system possesses (of course, for m degrees of freedom there are $(2m - 1)$ functionally independent constans of motion [21]); the arbitrary function C_1 has then a prescribed form [7], and μ is obtained by means of (20). Furthermore, it is possible to solve (11) in the one-dimensional case for μ, which yields a general solution in terms of Bessel functions. Using the initial conditions of the problem we find the particular solution. It is also possible to introduce the expression for the Lagrangean in the Euler-Lagrange equation and compare with the equation of motion to obtain μ[29]. We shall ilustrate here the first way.

It is not difficult to show that

$$C_2 \equiv \frac{1}{2} \dot{q}^2 - \frac{1}{2} k^2 q^2 \left(B - q \right)^2 = 0 \tag{21}$$

and a possible choice for C_1 is (see reference [7])

$$C_1 = c C_2^2 \tag{22}$$

where c is an arbitrary constant that multiplies the equation of motion. The Lagrangean reads then

$$\widetilde{L_V} = cC_2^2 \dot{q} \left[\ddot{q} - k^2 q (B-q)(B-2q) \right] \tag{23}$$

When inserted into the Euler-Lagrange equation, (23) provides the corresponding equation of motion (16) plus terms that are zero in virtue of the constant of motion. However, it has been shown that for the one-dimensional problem there exists an infinite number of Lagrangeans [8, 10]. Thus, we can write a more simpler s-equivalent one by means of the total time derivative of a certain gauge. Given the constant of motion, let choose

$$\frac{d\Lambda}{dt} = \dot{q}^2 - \left(C_1 \frac{dC_2}{dt} + C_2 \right) \tag{24}$$

Without loss of generality we set $c = \eta = 1$, and together with (24), equations (7) and (23) provide

$$L_V = \frac{1}{2}\dot{q}^2 + \frac{1}{2}k^2 q^2 \left(B-q\right)^2 \tag{25}$$

and the Hamiltonian can be written as

$$H_V = \frac{1}{2}P^2 - \frac{1}{2}k^2 q^2 \left(B-q\right)^2 \tag{26}$$

where the generalized momentum is

$$P \equiv \frac{\partial L}{\partial \dot{q}} = \dot{q} \tag{27}$$

IV. CONSERVATION, ENERGIES AND EQUILIBRIUM CONDITIONS

Now we proceed to study some of the applications the Lagrangean formalism offers to the Verhulst system.

The first term in (25) and (26) is identified as the usual traslational "kinetic" energy: quadratic and homogeneous in the first temporal derivative of the generalized coordinate. The second term on the right side of equation (26) is explicitily time-independent, meaning that H_V coincides with the energy function and it is then possible to write the potential energy of the system as

$$V = -\frac{1}{2}k^2 q^2 \left(B-q\right)^2 \tag{28}$$

The behavior of this potential is qualitatively presented in Figure 2 for fixed carrying capacity. In this potential, the smaller the time scale $(1/A)$, the more closed the curve and the larger the depression. Likewise, the greater the carrying capacity, B, the more pronounced the depression and the more open the curve. Equation (28) can be interpreted as an impulsor-retardatrice potential, depending only on the generalized coordinate. In fact, observe that when $q_0 \leq B$, for the interval $(0, B/2)$ the acceleration associated impulses the movement, being conversely for the interval $(B/2, B)$; the point $B/2$ corresponds to a local minimum, where the acceleration instantaneously annuls itself. On the other hand, when $q_0 > B$, i.e. the interval (B, ∞), the character of the acceleration is always impulsive, even being $\dot{q} < 0$. All this is consistent with the profiles sketched in Figure 1 and with the standard knowledge about equation (14).

Despite the fact the VLE describes multiple phenomena which one should think presents dissipation of energy (for example as in population dynamics, growth of living beings or meteorological models), it is easy to show that the Verhulst system is conservative (population systems with a first integral has been studied, see [12] and references therein); thus, the energy function H_V equals the total energy E. Note that in virtue of (14) equation (26) is identically zero. Taking all these considerations into account, it is clear in Figure 2 that the kinetic energy tends to annul itself as q approachs B. As a consequence, the system takes an infinite time to reach the steady state, and $q_0 < B$ implies always $q < B$ (similarly for $q_0 > B$).

Another interesting issue is that there are not oscillatory behaviours, despite the similarity with harmonic or the anharmonic potentials. Observe that as far as we are just interested in feassible results, we must take only the positive values of $q(t)$. Thus the character of the force (see equation (16)) is not restorative, but the commented before for each interval.

One important application is that of finding the stability cases. The usual derivative criteria allows us to find the equilibrium points, for which

$$\frac{\partial E}{\partial q} = 0 \tag{29}$$

The stability is then ruled by the condition

$$\frac{\partial^2 E}{\partial q^2} > 0 \tag{30}$$

The equilibrium solutions are $q_e = \left\{0, \frac{B}{2}, B\right\}$. The first one corresponds to a point of instability, the second to minimal potential energy and the last one is, as expected, the state of stationary equilibrium. As before, this analysis is in accordance with the known behaviour of the VLE (see Figure 1). It is important to remark here that usually the study of its equilibrium points is treated by means of a Taylor expansion of (14) about $q = 0$ and $q = B$ (see for example [12]). The variational formalism provides thus both a more elegant and complete treatment.

On the other hand, as far as the VLE can model various self-regulated phenomena, it should be useful for the description of the related statistical systems to write the total mean energy. We shall use the virial theorem for this porpouse. In our case,

$$\begin{aligned} \langle T \rangle &= -\frac{1}{2}\left\langle \frac{\partial V}{\partial q} q \right\rangle \\ &= \frac{1}{2} k^2 \left\langle q^2 (B-q)(B-2q) \right\rangle \end{aligned} \tag{31}$$

Then, the total mean energy is written as

$$\begin{aligned} \langle E \rangle \equiv \langle T + V \rangle &= -\frac{1}{2} k^2 \left\langle q^3 (B-q) \right\rangle \\ &= -\frac{k}{6}\left\langle \frac{d}{dt}\left(q^3\right) \right\rangle \end{aligned} \tag{32}$$

or, explicitily, evaluating the integral between $t = 0$ and $t \to \infty$,

$$\begin{aligned} \langle E \rangle_\infty &= \lim_{T\to\infty} \left\{ -\frac{k}{6T}\left[q(T)^3 - q_0^3 \right] \right\} \\ &= 0 \end{aligned} \tag{33}$$

Thus the total mean energy of the systems described by VLE is always null. But if as superior temporal bound we choose the characteristic time of the system, then

$$\langle E \rangle_{1/A} = -\frac{A^2}{6B}\left[q\left(\frac{1}{A}\right)^3 - q_0^3 \right] \tag{34}$$

Returning to the general case, observe that under certain circumstances equation (32) make it possible to stablish a relationship between the mean-squared velocity and the mean-linear velocity. Indeed, solving (14) for q we find two solutions:

$$q_\pm(t) = \frac{kB \pm \sqrt{B^2k^2 - 4k\dot{q}}}{2k} \tag{35}$$

Using $q_+(t)$, equation (32) can be restated as

$$\langle E \rangle = \left\langle \frac{\dot{q}^2}{2} - \frac{B}{4}\dot{q}\left[Bk + \sqrt{B^2k^2 - 4k\dot{q}} \right] \right\rangle \tag{36}$$

Then

$$\left\langle \dot{q}^2 \right\rangle = \frac{B}{2}\left\langle \dot{q}\left[Bk + \sqrt{B^2k^2 - 4k\dot{q}} \right] \right\rangle + 2\langle E \rangle \tag{37}$$

Note that if $\left| \frac{4\dot{q}}{B^2k} \right| \ll 1$ the latter equation simplifies to

$$\left\langle \dot{q}^2 \right\rangle = \frac{AB}{2}\langle \dot{q} \rangle + 2\langle E \rangle \tag{38}$$

The use of the other solution, $q_-(t)$, provides formally

$$\langle E \rangle = 0 \tag{39}$$

and then equation (37) is no longer true.

V. CONCLUDING REMARKS

We suggest in this work the use of the IPVC to obtain a variational formulation for self-limiting systems. This approach provides at least a Lagrangean, if it exists, departing from the equation of motion and regardless the system being conservative (when the definition $L = T - V$ holds) or not. There are several advantages, namely, it is possible

(a) to have in just one relation all the dynamical information of the system;

(b) to use a well know mathematical machinery to obtain from the Lagrangean or Hamiltonian the conserved quantities (and symmetries), equilibria points and trajectories, etc.;

(c) to group self-regulated phenomena having the same variational formulation, even when describing quite different scenarios, classifying the systems by the form of the Lagrangean or by its dynamical properties;

(d) to solve, when possible, the equation of motion. For some systems (as those described by Lotka-Volterra models) the Hamilton-Jacobi formalism, for example, should be useful to integrate the equation of motion;

(e) to help to understand how the models are related. Some models are, indeed, particular cases of more general equations;

(f) to test if a certain model is acceptable in terms of the Least-Action Principle.

On the other hand, making use of the Hojman et al. method, we have found the Lagrangean (25) and Hamiltonian (26) corresponding to the (1+1)-dimensional logistic equation as an example of the way a variational formalism can be obtained for general self-regulated systems. This may be regarded as a proof that the VLE corresponds to an extremal trajectory, thus it is physically acceptable (it complies with the Hamilton's Least-Action Principle) despite the fact Verhulst introduced it heuristically.

The potential energy (28) was briefly analized and the equilibria points and the stable/unstable cases were obtained by means of the usual derivative criteria, in accord with the other method (see [12]). Also, using the virial theorem it was possible to write an expression for the total mean energy of the system, equation (32), and a relation between the mean-squared velocity and the mean-linear velocity, equation (37).

It is not surprising to note that the Malthusian limit is trivially obtained for the particular case when B tends to ∞. Thus, the potential energy for this case can be written as

$$V_{Malthus} = -\frac{1}{2}A^2q^2 \tag{40}$$

We have in this work studied the thus-called second order approach. There is, certainly, a first order method of the Hojman et al. proceedure [7, 10]; we show briefly in the Appendix the way an s-equivalent Lagrangean for the VLE can be constructed using this approach. Even though the Lagrangean reproduces the logistic equation, as required, the physical interpretation of its terms is not so clear. However, generally speaking, it is important to bear in mind this proceedure because it sometimes provides a variational formalism of the physical phenomenon while the second order approach does not [7].

Finally, it could be useful to study in future works generalized auto-regulated models as for example those reported in references [23, 24, 25, 26, 27]. It is also interesting to extend this work to the study of the quantum systems related with self-regulated phenomena [28].

Acknowledgments

The authors wish to thank A. Rincón, A. Skirzewski, J.L. Flores, J. Toro-Mendoza, R.O. Rodríguez and D. Montiel for helpful comments. ÁGMS aknowlegde a Centro de Estudios de Postgrado (CEP) of the Universidad de Los Andes postgraduate student research grant. This work was partially supported by División de Investigación, Facultad de Ciencias, La Universidad del Zulia.

APPENDIX

In order to obtain an s-equivalent Lagrangean for the Verhulst's equation using the first order approach of the Hojman et al. method we proceed as follows.

First, let write the logistic equation as

$$\dot{q} = kq\,(B - q) \tag{41}$$

where $q \equiv q(t)$ is the generalized coordinate. Then, let write and choose a new integral variable $Q(t)$

$$Q(t) \equiv \int q(t)dt \tag{42}$$

such that equation (41) transforms in

$$\ddot{Q}(t) = k\dot{Q}(t)\left[B - \dot{Q}(t)\right] \tag{43}$$

This second order equation can be casted in two first order equations

$$\left\{ \begin{array}{c} \dot{x_1} = x_2 \\ \dot{x_2} = kx_2(B - x_2) \end{array} \right. \tag{44}$$

where $x_1 \equiv x_1(t)$ and $x_2 \equiv x_2(t)$. Note that indeed x_2 is exactly the generalized coordinate q.

The solution of this system of equations is given by

$$x_1 = \int x_2 dt + C_1 \tag{45}$$

$$x_2 = \frac{B}{1 + BC_2e^{-kBt}} \tag{46}$$

and this functional form for x_2 is just another way to write the usual form (2). Now, solving (45) and (46) for C_1 and C_2 we obtain

$$\begin{aligned} C_1 &= kx_1 - kBt - \ln B - \ln x_2 & (47) \\ C_2 &= \frac{e^{kBt}}{x_2}\left(1 - \frac{x_2}{B}\right) & (48) \end{aligned}$$

and following Hojman and Urrutia [7] the s-equivalent Lagrangean is finally written as

$$\begin{aligned} L \;=\; e^{kBt} \Bigg[& \frac{k\dot{x}_1}{x_2} - \frac{k\dot{x}_1}{B} - \frac{kB}{x_2} + \frac{\dot{x}_2}{x_2^2} - \frac{\dot{x}_2}{Bx_2} + \\ & - \frac{kB\ln(x_2)}{x_2} + k^2x_1 + k\ln n(x_2) + \frac{k\dot{x}_2x_1}{x_2^2} + \\ & - \frac{k^2Bx_1}{x_2} - \frac{k^2B^2t}{x_2} + \frac{kB\ln(B)}{x_2} + \\ & - \frac{kBt\dot{x}_2}{x_2^2} - \frac{\ln(B)\dot{x}_2}{x_2^2} + \frac{\ln(x_2)\dot{x}_2}{x_2^2} \Bigg] \end{aligned} \tag{49}$$

[1] Ván, P. & Nyíri, B., *Ann. Phys. (Leipzig)*, **8**, 4, 331-354, 1999.

[2] Vainberg, M.M., Methods for the Study of Nonlinear Operators, (Holden-Day, San Francisco, California, 1964).

[3] Tonti, E., *Int. Jour. Engin. Sci.,* **22**, 1343, 1984.

[4] Ichiyanagi, M., *Phys. Rep.,***243/3**, 125, 1994.

[5] Ván, P. & Muschik, W., *Phys. Rev. E*, **5/4**, 3584, 1995.

[6] Nyíri, B., *Journal of Non-Equilibrium Thermodynamics,* **16**, 217, 1991.

[7] Hojman, S., Urrutia, L.F., *J. Math. Phys.,* **22**, 1896, 1981.

[8] Hojman, S., Shepley, L.C., *Rev.Mex.Fís.*, **28**, 149, 1982.

[9] Helmholtz, H., *Journal für die reine und angewandte Mathematik* (Berlin), **100**, 137, 1887.

[10] Hojman, R., et al., *Phys. Rev. D.,***28**, 6, 1333-1336, 1983.

[11] Verhulst, P.F., *Corr. Math.et Phys.*, **10**, 113-121, 1838.

[12] Murray, J.D., Mathematical Biology, 2nd Edition, (Springer Verlag, Berlin, 1993).

[13] Malthus, T.R., An essay on the Principal of Population, 1798, (Penguin Books, 1970).

[14] Thompson, D.W., On Growth and Form, Revised Edition, Dover Pub., New York, 1992.

[15] Alonso, M. and Finn, E.J., Fundamental University Physics, Volume III, Quantum and statistical Physics (Addison-Wesley Publishing Company, Reading, Massachusetts, 1968), p. 344-345.

[16] Nelson, D.R. et al., *Physical Review E*, **58**, No. 2, 1383-1403, 1998.

[17] Euler, L., *Acta Acad. Sci. Imp. Petropol.*, 163-193, 1778.

[18] Townsend, J.S., Electricity in Gases (Oxford University Press, London, 1914).

[19] Joos, G., Theoretical Physics (Dover Publications, Mineola, New York, 1986), pp. 458-462.

[20] Pardo, F., *J. Math. Phys.,* **30**, No. 9, 2054-2061, 1989.

[21] Landau, L. , Mecanica, (Ed. Mir, Rome, 1975), p. 42.

[22] Goldstein, H., Classical Mechanics, 2nd Edition, (Adison-Wesley, London, 1980), p. 411.

[23] Foerster, Von H., et al., *Science*, **132**, 1291, 1960.

[24] Hoerner, von S.J., *British Interplanetary Society*, **28**, 691, 1975.

[25] Kapitza, S.P., *Uspehi Fisicheskjh Nauk (Russia)*, **166**, 1, 63-79, 1966.

[26] Kapitza, S.P., *How Many People Lived, Live and are to Live in the World. An essay on theory of growth of Humankind,* Moscow, Inst. Phys. Problem RAS, 1999.

[27] Kobelev, L. Ya., et al., *Will the Population of Humanity in the Future be Stabilized?*, http://arXiv.org/abs/physics/0003035, 16 Mar 2000.

[28] Muñoz S., A.G., Sierra P., D., Soldovieri, T., Rodriguez, O., *Quantization of Self-Regulated Systems* (in preparation), 2003.

[29] Even when not indicated by Hojman and colaborators, this idea may have some advantages. An adequate choice of the gauge and the use of equation (19) and (7) can simplify sometimes the calculations and the final Lagrangean.

Figure captions.

1. The Verhulst logistic equation for fixed carrying capacity ($B = 20$).

2. The Verhulst potential for differents k ($B = 5$).


40
35
30
25
20
15
10
5
0
0
0.5
1
1.5
2
2.5
3
3.5
4
4.5
t


0
-2000
-4000
-6000
-8000
-10000
V(q)
0
1
2
3
4
5
6
7
8
q
k=5
k=10
k=15
k=20